BLACK RIVER AGRICULTURAL NPS PLAN AND TMDL STRATEGY

Lorain and Medina Soil and Water Conservation Districts and Natural Resource Conservation Service December, 2007

1. RIPARIAN COORIDORS / STREAMS

Use existing USDA programs (CRP, CREP, EQIP, and WHIP) to improve stream corridors. Installing vegetative filter-strips of grass and/or trees in existing agricultural fields is the recommended method to improve water quality, treat erosion and create wildlife habitat.

Strategy:

- Establish grass / trees filter strip on 50% of agricultural field in watershed with in 10 years.
- Focus on educational programs to inform agricultural producers of filter strip programs and the benefits.

2. STREAMBANK EROSION

A. Removal of logiams and clearing / snagging is needed to facilitate flow and reduce the risk of stream-bank erosion.

Strategy:

- Develop a demonstration project, to remove logiams and major stream obstruction.
- B. Stream-bank erosion is a severe problem through most major channels within the watershed. What is the best method to deal with this problem?

Strategy:

 Use educational programs to inform landowners of methods of streambank erosion control.

3. LIVESTOCK

Provide financial and technical assistance to traditional and non-traditional livestock operations to manage waste and improve wastewater.

A. Conventional livestock operation (dairy, beef, and hogs)

Strategy

- Have all major livestock operations develop Comprehensive Nutrient Management Plan (CNMP) and or Grazing Management Plan to assist them with managing waste.
- Inventory livestock operations with animals grazing stream corridors.
- Provide Financial and technical assistance to all grazing operations to fence livestock from streams.
- Develop information program to inform livestock operators about rules concerning winter application of manure.
- Provide funding to livestock operations to assist with managing waste.
 Utilize EQIP and APAP,
- B. Non-Conventional livestock (horses, alpaca and others) operations. Provide technical assistance to non-traditional livestock operations to assist with waste management to improve water quality.

Strategy:

- Inventory non-traditional livestock operations.
- Education non-traditional livestock operators about state pollution abatement laws. (LEAP).
- Develop low-cost waste storage plans for use by non-traditional livestock operators.

4. AGRICULTURAL CROP PRODUCTION

Sheet and rill erosion from agricultural lands results in pollution from sedimentation and nutrients. This use of Best management Practices (BMP) is the recommended method to improve water quality.

Strategy:

- Encourage the use of permanent conservation practices (grass waterways, wascobs, erosion control structures) conservation tillage, filter strips, nutrient management as the primary BMP's to control erosion manage fertility and improve water quality.
- Utilize controlled drainage on tillable land to increase the effectiveness of conservation tillage and reduce nutrients and sedimentation to streams.

Utilize cover crops on low-residue crop fields (soybeans, corn sillage) to control erosion.

5. CHANNELIZATION

Ditch cleaning has traditionally been a double-edged sword. Residents feel it is needed to control localized flooding. Environmentalist discourage it because of it's impact to stream biology and wildlife. Local SWCD's are attempting to take a more balance approach to these opposing viewpoints when conducting drainage activities. They advocate limited drainage ditch improvement that mitigates local flooding issues without complete disruption to the stream corridor and surrounding habitat. This would include low-impact technique such as one-sided construction, maintaining natural vegetation on one bank, seeding of wildlife grasses and maintaining stream pools (were practical). This type of approach is the most practical solution to this complicated issue.